

## Well-Posedness of Parabolic Differential and Difference Equations with the Fractional Differential Operator

<sup>1,2</sup>Allaberen Ashyralyev

<sup>1</sup>*Department of Mathematics, ITTU, Ashgabat, Turkmenistan*

<sup>2</sup>*Department of Mathematics, Fatih University, Istanbul, Turkey*

*E-mail: aashyr@fatih.edu.tr*

### ABSTRACT

The stable difference scheme for the approximate solution of the initial value problem

$$\frac{du(t)}{dt} + D_t^{\frac{1}{2}}u(t) + Au(t) = f(t), \quad 0 < t < 1, \quad u(0) = 0$$
 for the differential equation in a

Banach space  $E$  with the strongly positive operator  $A$  and fractional operator  $D_t^{\frac{1}{2}}$  is presented. The well-posedness of the difference scheme in difference analogues of spaces of smooth functions is established. In practice, the coercive stability estimates for the solution of difference schemes for the  $2m$ -th order multi-dimensional fractional parabolic equation and the one-dimensional fractional parabolic equation with nonlocal boundary conditions in space variable are obtained.

2000 MSC: 65N12, 65M12

Keywords: Fractional parabolic equation, basset problems, well-posedness; coercive stability, difference scheme

### 1. INTRODUCTION

It is known that differential equations involving derivatives of non integer order have shown to be adequate models for various physical phenomena in areas like rheology, damping laws, diffusion processes, etc. (see the references given therein). A review of some applications of fractional calculus in continuum and statistical mechanics is given by Mainardi (1997).

The role played by coercive stability inequalities (well-posedness) in the study of boundary-value problems for parabolic partial differential and difference equations is well known (see Ashyralyev and Sobolevskii (1994, 2004)). In Ashyralyev (2010), the initial value problem

$$\frac{du(t)}{dt} + D_t^{\frac{1}{2}}u(t) + Au(t) = f(t), \quad 0 < t < 1, \quad u(0) = 0 \quad (1)$$

for the fractional differential equation in a Banach space  $E$  with the strongly positive operator  $A$  is considered. Here  $D_t^{\frac{1}{2}} = D_{0+}^{\frac{1}{2}}$  is the standard Riemann-Liouville's derivative of order  $\frac{1}{2}$ . This fractional differential equation corresponds to the Basset problem (Basset (1910)). It represents a classical problem in fluid dynamics where the unsteady motion of a particle accelerates in a viscous fluid due to the gravity of force.

A function  $u(t)$  is called a solution of the problem (1) if the following conditions are satisfied:

- (i)  $u(t)$  is continuously differentiable on the segment  $[0,1]$ ,
- (ii) The element  $u(t)$  belongs to  $D(A)$  for all  $t \in [0,1]$  and the function  $Au(t)$  is continuous on the segment  $[0,1]$ ,
- (iii)  $u(t)$  satisfies the equation and the initial condition (1).

A solution of problem (1) defined in this manner will from now on referred to as a solution of problem (1) in the space  $C(E) = C([0,1], E)$  of all continuous functions  $\varphi(t)$  defined on  $[0,1]$  with values in  $E$  equipped with the norm

$$\|\varphi\|_{C(E)} = \max_{0 \leq t \leq 1} \|\varphi(t)\|_E.$$

The well-posedness in  $C(E)$  of the boundary value problem (1) means that coercive inequality

$$\|u'\|_{C(E)} + \left\| D_t^{\frac{1}{2}} u \right\|_{C(E)} + \|Au\|_{C(E)} \leq M \|f\|_{C(E)}$$

is true for its solution  $u(t) \in C(E)$  with some  $M$ , which does not depend on  $f(t) \in C(E)$ .

Positive constants, which can differ in time (hence: not a subject of precision) will be indicated with an  $M$ . On the other hand  $M(\alpha, \beta, \dots)$  is used to focus on the fact that the constant depends only on  $\alpha, \beta, \dots$ .

In Ashyralyev (2010), the following theorems on well-posedness of (1) in spaces of smooth functions was established.

**Theorem 1.** Let  $A$  be a strongly positive operator in a Banach space  $E$  and  $f(t) \in C(E)$ . Then, for the solution  $u(t)$  in  $C(E)$  of the initial value problem (1) the stability inequality holds:

$$\left\| D_t^{\frac{1}{2}} u \right\|_{C(E)} + \|u' + Au\|_{C(E)} \leq M \|f\|_{C(E)}. \quad (2)$$

**Theorem 2.** Let  $A$  be a strongly positive operator in a Banach space  $E$  and  $f(t) \in C(E_\alpha)$  ( $0 < \alpha < 1$ ). Then for the solution  $u(t)$  in  $C(E_\alpha)$  of the initial value problem (1) the coercive inequality is valid:

$$\|u'\|_{C(E_\alpha)} + \|Au\|_{C(E_\alpha)} \leq M \alpha^{-1} (1 - \alpha)^{-1} \|f\|_{C(E_\alpha)}. \quad (3)$$

Here, the fractional space  $E_\alpha = E_\alpha(E, A)$  ( $0 < \alpha < 1$ ), consisting of all  $v \in E$  for which the following norm is finite:

$$\|v\|_{E_\alpha} = \sup_{\lambda > 0} \lambda^{1-\alpha} \|A \exp(-\lambda A)v\|_E$$

is additionally introduced.

In the present paper, the stable difference scheme for the approximate solution of initial value problem (1)

$$\begin{cases} \tau^{-1}(u_k - u_{k-1}) + Au_k + D_\tau^{\frac{1}{2}} u_k = f_k, \\ f_k - f(t_k), \quad t_k = k\tau, \quad 1 \leq k \leq N, \quad N\tau = 1, \quad u_0 = 0 \end{cases} \quad (4)$$

is presented. Here,

$$D_\tau^{\frac{1}{2}} u_k = \frac{1}{\sqrt{\pi}} \sum_{m=1}^k \frac{\Gamma(k-m+\frac{1}{2})}{(k-m)!} \frac{u_m - u_{m-1}}{\tau^{\frac{1}{2}}}, \quad \Gamma(k-m+\frac{1}{2}) = \int_0^\infty t^{k-m-\frac{1}{2}} e^{-t} dt. \quad (5)$$

The paper is organized as follows. The well-posedness of (4) in difference analogues of spaces of smooth functions is established in Section 2. In Section 3 the coercive stability estimates for the solution of difference schemes for the  $2m$ -th order multi-dimensional fractional parabolic equation and the one-dimensional fractional parabolic equation with nonlocal boundary conditions in space variable are obtained.

## 2. THE WELL-POSEDNESS OF DIFFERENCE SCHEME

Let us first obtain the representation for the solution of problem (4). It is clear that the first order of accuracy difference scheme

$$\tau^{-1}(u_k - u_{k-1}) + Au_k = F_k, \quad 1 \leq k \leq N\tau = 1, \quad u_0 = 0$$

has a solution and the following formula holds:

$$u_k = \sum_{s=1}^k R^{k-s+1} F_s \tau, \quad 1 \leq k \leq N,$$

where  $R = (I + \tau A)^{-1}$ . Applying the formula  $F_k = f_k - D_{\tau}^{\frac{1}{2}} u_k$ , we get

$$u_k = -\sum_{s=1}^k R^{k-s+1} D_{\tau}^{\frac{1}{2}} u_s \tau + \sum_{s=1}^k R^{k-s+1} f_s \tau, \quad 1 \leq k \leq N. \quad (6)$$

So, formula (6) gives the representation for the solution of problem (4). Let  $F_{\tau}(E)$  be the linear space of mesh functions  $\varphi^{\tau} = \{\varphi_k\}_1^N$  with values in the Banach space  $E$ . Next on  $F_{\tau}(E)$  we introduce the Banach space  $C_{\tau}(E) = C([0,1]_{\tau}, E)$  with the norm

$$\|\varphi^{\tau}\|_{C_{\tau}(E)} = \max_{1 \leq k \leq N} \|\varphi_k\|_E.$$

**Theorem 3.** Let  $A$  be a strongly positive operator in a Banach space  $E$ . Then, for the solution  $u^{\tau} = \{u_k\}_1^N$  in  $C_{\tau}(E)$  of initial value problem (4) the stability inequality holds:

$$\left\| \left\{ D_{\tau}^{\frac{1}{2}} u_k \right\}_1^N \right\|_{C_{\tau}(E)} + \left\| \left\{ \tau^{-1}(u_k - u_{k-1}) + Au_k \right\}_1^N \right\|_{C_{\tau}(E)} \leq M \|f^{\tau}\|_{C_{\tau}(E)}. \quad (7)$$

**Proof.** Using formula (6), we get

$$\tau^{-1}(u_k - u_{k-1}) = D_{\tau}^{\frac{1}{2}} u_k + \sum_{s=1}^k AR^{k-s+1} D_{\tau}^{\frac{1}{2}} u_s \tau + f_k - \sum_{s=1}^k AR^{k-s+1} f_s \tau. \quad (8)$$

Applying formulas (8) and (5), we obtain

$$\begin{aligned} D_{\tau}^{\frac{1}{2}} u_k &= \frac{1}{\sqrt{\pi}} \sum_{m=1}^k \frac{\Gamma(k-m+\frac{1}{2})}{(k-m)!} \tau^{\frac{1}{2}} \left[ -D_{\tau}^{\frac{1}{2}} u_m + f_m \right] \\ &+ \frac{1}{\sqrt{\pi}} \sum_{s=1}^k \sum_{m=s}^k \frac{\Gamma(k-m+\frac{1}{2})}{(k-m)!} AR^{m-s+1} D_{\tau}^{\frac{1}{2}} u_s \tau^{\frac{3}{2}} \\ &- \frac{1}{\sqrt{\pi}} \sum_{s=1}^k \sum_{m=s}^k \frac{\Gamma(k-m+\frac{1}{2})}{(k-m)!} AR^{m-s+1} f_s \tau^{\frac{3}{2}}. \end{aligned}$$

Let us first obtain the estimate

$$\left\| \frac{1}{\sqrt{\pi}} \sum_{m=s}^k \frac{\Gamma(k-m+\frac{1}{2})}{(k-m)!} AR^{m-s+1} \tau^{\frac{1}{2}} \right\|_{E \rightarrow E} \leq \frac{M}{\sqrt{(k-s)\tau}} \quad (9)$$

for any  $1 \leq s < k \leq N$ . We have that

$$\begin{aligned} \frac{1}{\sqrt{\pi}} \sum_{m=s}^k \frac{\Gamma(k-m+\frac{1}{2})}{(k-m)!} AR^{m-s+1} \tau^{\frac{1}{2}} &= \frac{1}{\sqrt{\pi}} \sum_{m=\lceil \frac{s+k}{2} \rceil}^k \frac{\Gamma(k-m+\frac{1}{2})}{(k-m)!} AR^{m-s+1} \tau^{\frac{1}{2}} \\ &+ \frac{1}{\sqrt{\pi}} \sum_{m=s}^{\lceil \frac{s+k}{2} \rceil} \frac{\Gamma(k-m+\frac{1}{2})}{(k-m)!} AR^{m-s+1} \tau^{\frac{1}{2}} = J_1 + J_2. \end{aligned}$$

Using estimates Sobolevskii (1971)

$$\|AR^k\|_{E \rightarrow E} \leq \frac{M}{k\tau}, \|R^k\|_{E \rightarrow E} \leq M, \quad 1 \leq k \leq N \quad (10)$$

and the following elementary inequality

$$\frac{\Gamma(k-m+\frac{1}{2})}{(k-m)!} \leq \frac{1}{\sqrt{k-m}}, \quad 0 \leq m < k, \quad (11)$$

we get

$$\begin{aligned} \|J_1\|_{E \rightarrow E} &\leq \frac{1}{\sqrt{\pi}} \sum_{m=\lfloor \frac{s+k}{2} \rfloor}^k \frac{\Gamma(k-m+\frac{1}{2})}{(k-m)!} \|AR^{m-s+1}\|_{E \rightarrow E} \tau^{\frac{1}{2}} \\ &\leq \frac{2M}{(k-s)\tau} \frac{1}{\sqrt{\pi}} \sum_{m=\lfloor \frac{s+k}{2} \rfloor}^k \frac{\tau}{\sqrt{(k-m)\tau}} \leq \frac{M_1}{\sqrt{(k-s)\tau}}. \end{aligned} \tag{12}$$

Now, we shall estimate  $J_2$ . We have that

$$\begin{aligned} J_2 &= \frac{1}{\sqrt{\pi}} \frac{\Gamma(k-s+\frac{1}{2})}{(k-s)!} \tau^{-\frac{1}{2}} - \frac{1}{\sqrt{\pi}} \frac{\Gamma(k-\lfloor \frac{s+k}{2} \rfloor + \frac{3}{2})}{(k-\lfloor \frac{s+k}{2} \rfloor + 1)!} R^{\lfloor \frac{s+k}{2} \rfloor - s} \tau^{-\frac{1}{2}} \\ &\quad + \frac{1}{\sqrt{\pi}} \sum_{m=s+1}^{\lfloor \frac{s+k}{2} \rfloor - 1} \left[ \frac{\Gamma(k-m+\frac{1}{2})}{(k-m)!} - \frac{\Gamma(k-m+\frac{3}{2})}{(k-m+1)!} \right] R^{m-s} \tau^{-\frac{1}{2}}. \end{aligned}$$

Applying estimates (10) and (11), we obtain

$$\begin{aligned} \|J_2\|_{E \rightarrow E} &\leq \frac{1}{\sqrt{\pi}} \frac{1}{\sqrt{(k-s)\tau}} + \frac{1}{\sqrt{\pi}} \|R^{\lfloor \frac{s+k}{2} \rfloor - s}\|_{E \rightarrow E} \frac{1}{\sqrt{(k-\lfloor \frac{s+k}{2} \rfloor + 1)\tau}} \\ &\quad + \frac{1}{\sqrt{\pi}} \sum_{m=s+1}^{\lfloor \frac{s+k}{2} \rfloor - 1} \left| \frac{\Gamma(k-m+\frac{1}{2})}{(k-m)!} - \frac{\Gamma(k-m+\frac{3}{2})}{(k-m+1)!} \right| \|R^{m-s}\|_{E \rightarrow E} \tau^{\frac{1}{2}} \\ &\leq \frac{1}{\sqrt{\pi}} \frac{1}{\sqrt{(k-s)\tau}} + \frac{1}{\sqrt{\pi}} M \frac{\sqrt{2}}{\sqrt{(k-s)\tau}} \\ &\quad + M \frac{1}{\sqrt{\pi}} \sum_{m=s+1}^{\lfloor \frac{s+k}{2} \rfloor - 1} \frac{\tau}{(k-m+1)\tau\sqrt{(k-m)\tau}} \leq \frac{M_2}{\sqrt{(k-s)\tau}}. \end{aligned} \tag{13}$$

Estimate (9) follows from estimates (12) and (13).

Now, let us first estimate  $z_k = \left\| D_{\tau}^{\frac{1}{2}} u_k \right\|_E$ . Applying the triangle inequality and estimate (9), we get

$$\begin{aligned} z_k &\leq \frac{1}{\sqrt{\pi}} \sum_{m=1}^k \frac{\Gamma(k-m+\frac{1}{2})}{(k-m)!} \tau^{-\frac{1}{2}} [z_m + \|f_m\|_E] \\ &\quad + \frac{1}{\sqrt{\pi}} \sum_{s=1}^k \left\| \sum_{m=s}^k \frac{\Gamma(k-m+\frac{1}{2})}{(k-m)!} AR^{m-s+1} \right\|_{E \rightarrow E} z_s \tau^{\frac{3}{2}} \\ &\quad + \frac{1}{\sqrt{\pi}} \sum_{s=1}^k \left\| \sum_{m=s}^k \frac{\Gamma(k-m+\frac{1}{2})}{(k-m)!} AR^{m-s+1} \right\|_{E \rightarrow E} \|f_s\|_E \tau^{\frac{3}{2}} \\ &\leq M_3 \sum_{s=1}^{k-1} \frac{1}{\sqrt{(k-s)\tau}} \tau [z_s + \|f_s\|_E] + M_4 [z_k + \|f_k\|_E] \tau^{\frac{1}{2}} \end{aligned}$$

for any  $k=1, \dots, N$ . Applying the above inequality and the difference analogue of the integral inequality, we obtain

$$\left\| \left\{ D_{\tau}^{\frac{1}{2}} u_k \right\}_1^N \right\|_{C_{\tau}(E)} \leq M \|f^{\tau}\|_{C_{\tau}(E)}. \tag{14}$$

Using the triangle inequality and equation (4), we get

$$\begin{aligned} \left\| \left\{ \tau^{-1}(u_k - u_{k-1}) + Au_k \right\}_1^N \right\|_{C_{\tau}(E)} &\leq \left[ \|f^{\tau}\|_{C_{\tau}(E)} + \left\| \left\{ D_{\tau}^{\frac{1}{2}} u_k \right\}_1^N \right\|_{C_{\tau}(E)} \right] \\ &\leq M_1 \|f^{\tau}\|_{C_{\tau}(E)}. \end{aligned} \tag{15}$$

Estimate (7) follows from estimates (14) and (15). Theorem 3 is proved.

**Theorem 4.** Let  $A$  be a strongly positive operator in a Banach space  $E$ . Then, for the solution  $u^\tau = \{u_k\}_1^N$  in  $C_\tau(E)$  of initial value problem (4) the almost coercive stability inequality is valid:

$$\begin{aligned} & \left\| \left\{ \tau^{-1}(u_k - u_{k-1}) \right\}_1^N \right\|_{C_\tau(E)} + \left\| \{Au_k\}_1^N \right\|_{C_\tau(E)} \\ & \leq M \min \left\{ \ln \frac{1}{\tau}, 1 + \ln \|A\|_{E \rightarrow E} \right\} \|f^\tau\|_{C_\tau(E)}. \end{aligned} \tag{16}$$

**Proof.** The proof of estimate

$$\left\| \left\{ \tau^{-1}(u_k - u_{k-1}) \right\}_1^N \right\|_{C_\tau(E)} \leq M \min \left\{ \ln \frac{1}{\tau}, 1 + \ln \|A\|_{E \rightarrow E} \right\} \|f^\tau\|_{C_\tau(E)} \tag{17}$$

or the solution of initial value problem (4) is based on estimate (7) and the following estimates (Ashyralyev and Sobolevskii (1994)):

$$\begin{aligned} \max_{1 \leq k \leq N} \left\| \sum_{s=1}^k AR^{k-s+1} f_s \tau \right\|_E & \leq M \min \left\{ \ln \frac{1}{\tau}, 1 + \ln \|A\|_{E \rightarrow E} \right\} \|f^\tau\|_{C(E)}, \\ \max_{1 \leq k \leq N} \left\| \sum_{s=1}^k AR^{k-s+1} D_\tau^{\frac{1}{2}} u_s \tau \right\|_E & \leq M \min \left\{ \ln \frac{1}{\tau}, 1 + \ln \|A\|_{E \rightarrow E} \right\} \left\| \left\{ D_\tau^{\frac{1}{2}} u_k \right\}_1^N \right\|_{C(E)}. \end{aligned}$$

Using these estimates, the triangle inequality and equation (4), we get

$$\left\| \{Au_k\}_1^N \right\|_{C_\tau(E)} \leq M \min \left\{ \ln \frac{1}{\tau}, 1 + \ln \|A\|_{E \rightarrow E} \right\} \|f^\tau\|_{C(E)}. \tag{18}$$

Estimate (16) follows from estimates (17) and (18). Theorem 4 is proved.

Note that the Banach space  $E'_\alpha = E'_\alpha(E, A)$  ( $0 < \alpha < 1$ ) consists of those  $v \in E$  for which the norm

$$\|v\|_{E'_\alpha} = \sup_{\lambda > 0} \lambda^\alpha \|A(\lambda + A)^{-1}v\|_E$$

is finite.

**Theorem 5.** Let  $A$  be a strongly positive operator in a Banach space  $E$ . Then, for the solution  $u^\tau = \{u_k\}_1^N$  in  $C_\tau(E'_\alpha)$  of the initial value problem (4) the coercive stability inequality is valid:

$$\begin{aligned} & \left\| \left\{ \tau^{-1}(u_k - u_{k-1}) \right\}_1^N \right\|_{C_\tau(E'_\alpha)} + \left\| \{Au_k\}_1^N \right\|_{C_\tau(E'_\alpha)} \\ & \leq M \alpha^{-1} (1 - \alpha)^{-1} \|f^\tau\|_{C_\tau(E'_\alpha)}. \end{aligned} \quad (19)$$

**Proof.** By Theorem 3,

$$\left\| \left\{ D_\tau^{\frac{1}{2}} u_k \right\}_1^N \right\|_{C_\tau(E'_\alpha)} \leq M \|f^\tau\|_{C_\tau(E'_\alpha)} \quad (20)$$

for the solution of initial value problem (4). The proof of estimate

$$\left\| \left\{ \tau^{-1}(u_k - u_{k-1}) \right\}_1^N \right\|_{C_\tau(E'_\alpha)} \leq M \alpha^{-1} (1 - \alpha)^{-1} \|f^\tau\|_{C_\tau(E'_\alpha)} \quad (21)$$

for the solution of initial value problem (4) is based on estimate (20) and the following estimates (Ashyralyev and Sobolevskii (1994)):

$$\max_{1 \leq k \leq N} \left\| \sum_{s=1}^k AR^{k-s+1} f_s \tau \right\|_{E'_\alpha} \leq M \alpha^{-1} (1 - \alpha)^{-1} \|f^\tau\|_{C_\tau(E'_\alpha)}, \quad (22)$$

$$\max_{1 \leq k \leq N} \left\| \sum_{s=1}^k AR^{k-s+1} D_\tau^{\frac{1}{2}} u_s \tau \right\|_{E'_\alpha} \leq M \alpha^{-1} (1 - \alpha)^{-1} \left\| \left\{ D_\tau^{\frac{1}{2}} u_k \right\}_1^N \right\|_{C_\tau(E'_\alpha)}. \quad (23)$$

Using the triangle inequality, estimates (22), (23) and equation (4), we get

$$\left\| \{Au_k\}_1^N \right\|_{C_\tau(E'_\alpha)} \leq M_1 \alpha^{-1} (1 - \alpha)^{-1} \|f^\tau\|_{C_\tau(E'_\alpha)}. \quad (24)$$

Estimate (19) follows from estimates (21) and (24). Theorem 5 is proved.

Note that by passing to the limit for  $\tau \rightarrow 0$  one can recover Theorems 1 and 2.

### 3. APPLICATIONS

Now, we consider the applications of Theorem 3, 4 and 5.

First, the initial-value problem on the range  $\{0 \leq t \leq 1, x \in \mathbb{R}^n\}$  for the  $2m$ -order multi-dimensional fractional parabolic equation is considered:

$$\begin{cases} \frac{\partial v(t, x)}{\partial t} + D_t^{\frac{1}{2}} v(t, x) + \sum_{|r|=2m} a_r(x) \frac{\partial^{|r|} v(t, x)}{\partial x_1^{r_1} \cdots \partial x_n^{r_n}} + \sigma v(t, x) = f(t, x), \\ 0 < t < 1; v(0, x) = 0, x \in \mathbb{R}^n, |r| = r_1 + \cdots + r_n, \end{cases} \quad (25)$$

where  $a_r(x)$  and  $f(t, x)$  are given as sufficiently smooth functions. Here,  $\sigma$  is a sufficiently large positive constant.

The discretization of problem (25) is carried out in two steps. In the first step, the grid space  $\mathbb{R}_h^n (0 < h \leq h_0)$  is defined as the set of all points of the Euclidean space  $\mathbb{R}^n$  whose coordinates are given by

$$x_k = s_k h, \quad s_k = 0, \pm 1, \pm 2, \dots, k = 1, \dots, n.$$

The difference operator  $A_h^x = B_h^x + \sigma I_h$  is assigned to the differential operator  $A^x = B^x + \sigma I$ , defined by (25). The operator

$$B_h^x = h^{-2m} \sum_{2m \leq |s| \leq S} b_s^x \Delta_{1-}^{s_1} \Delta_{1+}^{s_2} \cdots \Delta_{n-}^{s_{2n-1}} \Delta_{n+}^{s_{2n}}, \quad (26)$$

acts on functions defined on the entire space  $\mathbb{R}_h^n$ . Here,  $s \in \mathbb{R}^{2n}$  is a vector with nonnegative integer coordinates,

$$\Delta_{k\pm} f^h(x) = \pm (f^h(x \pm e_k h) - f^h(x)),$$

where  $e_k$  is the unit vector of the axis  $x_k$ .

An infinitely differentiable function  $\varphi(x)$  of the continuous argument  $x \in \mathbb{R}^n$  that is continuous and bounded together with all its derivatives is said to be smooth.

We say that the difference operator  $A_h^x$  is a  $\lambda$ -th order ( $\lambda > 0$ ) approximation of the differential operator  $A^x$  if the inequality

$$\sup_{x \in \mathbb{R}_h^n} |A_h^x \varphi(x) - A^x \varphi(x)| \leq M(\varphi)h^\lambda$$

holds for any smooth function  $\varphi(x)$ . The coefficients  $b_s^x$  are chosen in such a way that the operator  $A_h^x$  approximates in a specified way the operator  $A^x$ . It is assumed that the operator  $A_h^x$  approximates the differential operator  $A^x$  with any prescribed order (Smirnitskii and Sobolevskii (1981)).

The function  $A^x(\xi h, h)$  is obtained by replacing the operator  $\Delta_{k\pm}$  in the right-hand side of equality (26) with the expression  $\pm(\exp\{\pm i\xi_k h\} - 1)$ , respectively and is called the symbol of the difference operator  $B_h^x$ .

We shall assume that for  $|\xi_k h| \leq \pi$  and fixed  $x$  the symbol  $A^x(\xi h, h)$  of the operator  $B_h^x = A_h^x - \sigma I_h$  satisfies the inequalities

$$(-1)^m A^x(\xi h, h) \geq M |\xi|^{2m}, \left| \arg A^x(\xi h, h) \right| \leq \phi < \phi_0 \leq \frac{\pi}{2}. \quad (27)$$

Suppose that the coefficient  $b_s^x$  of the operator  $B_h^x = A_h^x - \sigma I_h$  is bounded and satisfies the inequalities

$$|b_s^{x+e_k h} - b_s^x| \leq Mh^\epsilon, x \in \mathbb{R}_h^n, \epsilon \in (0, 1]. \quad (28)$$

With the help of  $A_h^x$ , we arrive at the initial value problem

$$\begin{cases} \frac{\partial v(t, x)}{\partial t} + D_t^{\frac{1}{2}} v^h(t, x) + A_h^x v^h(t, x) = f^h(t, x), & 0 < t < 1, \quad x \in \mathbb{R}_h^n, \\ v^h(0, x) = 0, & x \in \mathbb{R}_h^n \end{cases} \quad (29)$$

for an infinite system of ordinary differential equations.

In the second step, problem (29) is replaced by the difference scheme

$$\begin{cases} \frac{u_k^h(x) - u_{k-1}^h(x)}{\tau} + D_\tau^{\frac{1}{2}} u_k^h + A_h^x u_k^h = f_k^h(x), \\ f_k^h(x) = f^h(t_k, x), \quad t_k = k\tau, \quad 1 \leq k \leq N, \quad N\tau = 1, \quad x \in \mathbb{R}_h^n, \\ u_0^h(x) = 0, \quad x \in \mathbb{R}_h^n. \end{cases} \quad (30)$$

Based on the number of corollaries of the abstract theorems given in the above, to formulate the result, one needs to introduce the spaces  $C_h = C(\mathbb{R}_h^n)$  and  $C_h^\beta = C^\beta(\mathbb{R}_h^n)$  of all bounded grid functions  $u^h(x)$  defined on  $\mathbb{R}_h^n$ , equipped with the norms

$$\|u^h\|_{C_h} = \sup_{x \in \mathbb{R}_h^n} |u^h(x)|, \quad \|u^h\|_{C_h^\beta} = \sup_{x \in \mathbb{R}_h^n} |u^h(x)| + \sup_{x, y \in \mathbb{R}_h^n} \frac{|u^h(x) - u^h(x+y)|}{|y|^\beta}.$$

**Theorem 6.** Suppose that assumptions (27) and (28) for the operator  $A_h^x$  hold. Then, the solutions of the difference scheme (30) satisfy the following stability estimates:

$$\begin{aligned} \max_{1 \leq k \leq N} \|D_\tau^{\frac{1}{2}} u_k^h\|_{C_h^\mu} &\leq M_1(\mu) \max_{1 \leq k \leq N} \|f_k^h\|_{C_h^\mu}, \quad 0 \leq \mu \leq 1, \\ \|\{\tau^{-1}(u_k^h - u_{k-1}^h)\}_1^N\|_{C_\tau(C_h^\mu)} &\leq M(\mu) \ln \frac{1}{\tau + |h|} \max_{1 \leq k \leq N} \|f_k^h\|_{C_h^\mu}, \quad 0 \leq \mu \leq 1, \\ \|\{\tau^{-1}(u_k^h - u_{k-1}^h)\}_1^N\|_{C_\tau(C_h^{\mu+2m\alpha})} &\leq M(\alpha, \mu) \max_{1 \leq k \leq N} \|f_k^h\|_{C_h^{\mu+2m\alpha}}, \quad 0 < 2m\alpha + \mu < 1. \end{aligned}$$

The proof of Theorem 5 is based on the abstract Theorems 3, 4 and 5, the strongly positivity of the operator  $A_h^x$  defined by (33) in  $C_h^\mu$  (Smiritskii and Sobolevskii (1981)) and the estimate

$$\min \left\{ \ln \frac{1}{\tau}, 1 + \ln \|A_h^x\|_{C_h^\mu \rightarrow C_h^\mu} \right\} \leq M(\mu) \ln \frac{1}{\tau + |h|}$$

and on the following two theorems on the coercivity inequality for the solution of the elliptic difference equation in  $C_h^\beta$  and on the structure of the fractional space  $E'_\alpha(C_h, A_h^x)$ .

**Theorem 7** (Ashyralyev and Sobolevskii (1994)). Suppose that assumptions (27) and (28) for the operator  $A_h^x$  hold.

Then, for the solution of the elliptic difference equation

$$A_h^x u^h(x) = \omega^h(x), x \in \mathbb{R}_h^n \tag{31}$$

the estimate

$$\sum_{2m \leq |s| \leq S} h^{-2m} \left\| \Delta_{1-}^{s_1} \Delta_{1+}^{s_2} \dots \Delta_{n-}^{s_{2n-1}} \Delta_{n+}^{s_{2n}} u^h \right\|_{C_h^\beta} \leq M(\sigma, \beta) \left\| \omega^h \right\|_{C_h^\beta}$$

is valid.

**Theorem 8** (Ashyralyev and Sobolevskii (1994)). Suppose that assumptions (27) and (28) for the operator  $A_h^x$  hold.

Then, for any  $0 < \alpha < \frac{1}{2m}$  the norms in the spaces  $E'_\alpha(C_h, A_h^x)$  and  $C_h^{2m\alpha}$  are equivalent uniformly in  $h$ .

Second, we consider the mixed boundary value problem for the fractional parabolic equation

$$\left\{ \begin{array}{l} \frac{\partial v(t, x)}{\partial t} D_t^{\frac{1}{2}} v(t, x) - a(x) \frac{\partial^2 v(t, x)}{\partial x^2} + \sigma v(t, x) = f(t, x), \\ 0 < t < 1, \quad 0 < x < 1; \quad v(0, x) = 0, \quad 0 \leq x \leq 1, \\ u(t, 0) = u(t, 1), \quad u_x(t, 0) = u_x(t, 1), \quad 0 \leq t \leq 1, \end{array} \right. \tag{32}$$

where  $a(t, x)$  and  $f(t, x)$  are given sufficiently smooth functions and  $a(t, x) \geq a > 0$ . Here,  $\sigma$  is a sufficiently large positive constant.

The discretization of problem (32) is carried out in two steps. In the first step, let us define the grid space

$$[0,1]_h = \{x: x_r = rh, 0 \leq r \leq K, Kh = 1\}.$$

We introduce the Banach space  $C_h^\beta = C^\beta([0,1]_h)$  ( $0 < \beta < 1$ ) of the grid functions  $\varphi^h(x) = \{\varphi_r\}_1^{K-1}$  defined on  $[0,1]_h$ , equipped with the norm

$$\|\varphi^h\|_{C_h^\beta} = \|\varphi^h\|_{C_h} + \sup_{1 \leq k < k+\tau \leq K-1} \frac{|\varphi_{k+\tau} - \varphi_k|}{\tau^\beta},$$

where  $C_h = C([0,1]_h)$  is the space of the grid functions defined on  $[0,1]_h$ , equipped with the norm

$$\|\varphi^h\|_{C_h} = \max_{1 \leq k \leq K-1} |\varphi_k|.$$

To the differential operator  $A$  generated by the problem (32), we assign the difference operator  $A_h^x$  by the formula

$$A_h^x \varphi^h(x) = \left\{ -\left(a(x)\varphi_x\right)_{x,r} + \delta\varphi_r \right\}_1^{K-1}, \quad (33)$$

acting in the space of grid functions  $\varphi^h(x) = \{\varphi_r\}_0^K$  satisfying the conditions  $\varphi_0 = \varphi_K, \varphi_1 - \varphi_0 = \varphi_K - \varphi_{K-1}$ . With the help of  $A_h^x$ , we arrive at the initial value problem

$$\begin{cases} \frac{d\varphi^h(t,x)}{dt} + D_t^{\frac{1}{2}} \varphi^h(t,x) + A_h^x \varphi^h(t,x) = f^h(t,x), & 0 < t < 1, \quad x \in [0,1]_h, \\ \varphi^h(0,x) = 0, & x \in [0,1]_h \end{cases} \quad (34)$$

for an infinite system of ordinary fractional differential equations. In the second step, we replace problem (34) by difference scheme (4)

$$\begin{cases} \frac{u_k^h(x) - u_{k-1}^h(x)}{\tau} + D_{\tau}^{\frac{1}{2}} u_k^h + A_h^x u_k^h(x) = f_k^h(x), & f_k^h(x) = \{f(t_k, x_r)\}_1^{K-1}, \\ t_k = k\tau, 1 \leq k \leq N, \quad N\tau = 1; \quad u_0^h(x) = 0, \quad x \in [0, 1]_h. \end{cases} \quad (35)$$

**Theorem 9.** Let  $\tau$  and  $h$  be sufficiently small numbers. Then, the solutions of the difference scheme (35) satisfy the following stability estimates

$$\begin{aligned} \max_{1 \leq k \leq N} \left\| D_{\tau}^{\frac{1}{2}} u_k^h \right\|_{C_h^{\mu}} &\leq M_1(\mu) \max_{1 \leq k \leq N} \|f_k^h\|_{C_h^{\mu}}, \quad 0 \leq \mu \leq 1, \\ \left\| \left\{ \tau^{-1}(u_k^h - u_{k-1}^h) \right\}_1^N \right\|_{C_{\tau}(C_h^{\mu})} &\leq M(\mu) \ln \frac{1}{\tau + |h|} \max_{1 \leq k \leq N} \|f_k^h\|_{C_h^{\mu}}, \quad 0 \leq \mu \leq 1, \\ \left\| \left\{ \tau^{-1}(u_k^h - u_{k-1}^h) \right\}_1^N \right\|_{C_{\tau}(C_h^{\mu+2\alpha})} &\leq M(\alpha, \mu) \max_{1 \leq k \leq N} \|f_k^h\|_{C_h^{\mu+2\alpha}}, \quad 0 < 2\alpha + \mu < 1. \end{aligned}$$

The proof of Theorem 9 is based on the abstract Theorems 3, 4 and 5, the strongly positivity of the operator  $A_h^x$  defined by (33) in  $C_h^{\mu}$  and the estimate

$$\min \left\{ \ln \frac{1}{\tau}, 1 + \ln \|A_h^x\|_{C_h^{\mu} \rightarrow C_h^{\mu}} \right\} \leq M(\mu) \ln \frac{1}{\tau + h}$$

and on the following theorem on the structure of the fractional space  $E'_{\alpha}(C_h, A_h^x)$ .

**Theorem 10** (Ashyralyev (2007)). For any  $0 < \alpha < \frac{1}{2}$ , the norms in the spaces  $E'_{\alpha}(C_h, A_h^x)$  and  $C_h^{2\alpha}$  are equivalent uniformly in  $h$ .

## REFERENCES

- Ashyralyev, A. 2010. Well-posedness of the Basset problem in spaces of smooth functions. *Applied Mathematics Letters*, 6 pages.
- Ashyralyev, A. 2009. A note on fractional derivatives and fractional powers of operators. *Journal of Mathematical Analysis and Applications*. **357**(1): 232-236.

- Ashyralyev, A., Dal, F. and Pinar, Z. 2009. On the numerical solution of fractional hyperbolic partial differential equations, *Mathematical Problems in Engineering*, Article ID 730465, 2009: 11 pages.
- Ashyralyev, A. 2007. Fractional spaces generated by the positive differential and difference operator in a Banach space. In: Kenan Tas *et al.*(ed) *Proc. of the Conf. Mathematical Methods and Engineering*. Springer, Netherlands, 13-22.
- Ashyralyev, A. and Sobolevskii, P. E. 2004. *New Difference Schemes for Partial Differential Equations*, Operator Theory: Advances and Applications. Basel, Boston, Berlin: Birkh user Verlag.
- Ashyralyev, A. and Sobolevskii, P.E. 1994. *Well-Posedness of Parabolic Difference Equations*, Operator Theory: Advances and Applications. Basel, Boston, Berlin: Birkhauser Verlag.
- Bagley, R. L. and Torvik, P. J. 1984. On the appearance of the fractional derivative in the behavior of real materials. *J. Appl. Mech.* **51**: 294-298.
- Basset, A.B. 1910. On the descent of a sphere in a viscous liquid. *Quart. J. Math.* **42**: 369-381.
- El-Mesiry, E.M., El-Sayed, A. M. A. and El-Saka, H. A. A. 2005. Numerical methods for multi-term fractional (arbitrary) orders differential equations. *Appl. Math. Comput.* **160**(3): 683-699.
- Lavoie, J. L., Osler, T. J. and Tremblay, R. 1976. Fractional derivatives and special functions. *SIAM Review.* **18**(2): 240-268.
- Mainardi, F. 1997. Fractional calculus: Some basic problems in continuum and statistical mechanics. In: A. Carpinteri and F. Mainardi, Ed., *Fractals and Fractional Calculus in Continuum Mechanics*, Springer-Verlag, New York: 291-348.
- Podlubny, I. 1999. *Fractional Differential Equations*. New York: Academic Press.

- Samko, S. G., Kilbas, A. A. and Marichev, O. I. 1993. *Fractional Integrals and Derivatives*. London: Gordon and Breach Science Publishers.
- Shaher Momani and Kamel Al-Khaled. 2005. Numerical solutions for systems of fractional differential equations by the decomposition method. *Applied Mathematics and Computation*. **162**(3): 1351-1365.
- Smirnitskii, Yu. A. and Sobolevskii, P. E. 1981. Positivity of multidimensional difference operators in the  $C$ -norm. *Usp. Mat. Nauk*. **36**(4): 202-203. (Russian).
- Sobolevskii, P. E. 1971. The coercive solvability of difference equations. *Dokl. Acad. Nauk SSSR*. **201**(5):1063-1066. (Russian).
- Tarasov, V. E. 2007. Fractional derivative as fractional power of derivative. *International Journal of Mathematics*. **18**: 281-299.